arXiv:cond-mat/9812248v1 [cond-mat.str-el] 15 Dec 1998

The spin gap of CaV_4O_9 revisited

M. Mambrini and F. Mila

Laboratoire de Physique Quantique, Université Paul Sabatier, 118 Route de Narbonne, 31062 Toulouse Cedex, France.

The large-plaquette scenario of the spin gap in CaV_4O_9 is investigated on the basis of extensive exact diagonalizations. We confirm the existence of a large-plaquette phase in a wide range of parameters, and we show that the most recent neutron scattering data actually *require* an intra-plaquette second neighbor exchange integral much larger than the inter-plaquette one, thus justifying the perturbative calculation used in the interpretation of the neutron scattering experiments.

PACS Nos : 75.10.Jm 75.40.Cx 75.50.Ee

CaV_4O_9 is the only known two-dimensional magnet presenting a gap in the spin excitations. The low temperature dependence of the magnetic susceptibility first measured by Taniguchi *et al.* [1] provided a value of $\Delta \approx 107$ K. Since then several scenarios based on the Heisenberg model on the depleted lattice have been proposed to explain both the presence and the value of Δ. It appears that the presence of frustration induced by a second neighbor antiferromagnetic interaction is necessary and it has become clear that two types of first and second neighbors exchange integrals are required to fit experiments correctly. The Hamiltonian then reads,

$$\mathcal{H} = J_1 \sum_{\langle i,j\rangle_p} \mathbf{S}_i.\mathbf{S}_j + J_1' \sum_{\langle i,j\rangle_d} \mathbf{S}_i.\mathbf{S}_j + J_2 \sum_{\langle\langle i,j\rangle\rangle_p} \mathbf{S}_i.\mathbf{S}_j + J_2' \sum_{\langle\langle i,j\rangle\rangle_d} \mathbf{S}_i.\mathbf{S}_j \qquad (1)$$

where the sums run respectively over bonds within plaquettes (p) and dimer between plaquettes (d) for first and second nearest neighbor.

The first explanation proposed to explain the presence of a gap was based on a $J_1 - J_2$ model on the depleted lattice with $J_2/J_1 = 1/2$ leading to weakly coupled small plaquettes. It must be abandoned for at least two reasons: on one hand, recent estimates of the exchange integrals in other members of the vanadates family suggest that the ratio J_2/J_1 is much larger than 1/2. On the other hand, the most recent data obtained with inelastic neutron scattering seem to be inconsistent with the picture of weakly coupled plaquettes built on the smallest squares of the depleted lattice. In particular, the dispersion curve of the lowest triplet state measured by Kodama *et al.* [2] turns out to be minimum at $(0,0)$ momentum. The authors then performed fits of the curve assuming that one of the integral exchange is large enough to justify a second order perturbation theory in the three other parameters and concluded that the best set of exchange integral is the one corresponding to J_1, J_1', J_2' small compared with J_2, namely $J_1 = J_1' \approx 67$ K, $J_2 \approx 171$ K and $J_2' \approx 15$ K. These parameters are also consistent with the temperature dependence of the susceptibility if one assumes that part of the sample is non-magnetic [3]. In this picture the system consists of a set of weakly coupled J_2-plaquettes. Nevertheless, regarding the mechanism responsible for the values of J_2 and J_2' (super-exchange between two second neighbor Vanadium atoms *via* an oxygen atom), the fact that these integrals differ by one order of magnitude is *a priori* surprising, although it is not necessarily contradictory since exchange integrals are very difficult to evaluate directly (from *ab initio* method in quantum chemistry for example).



FIG. 1. Symmetry of the first excited state for $J_1' = J_1$. Symmetry sectors are labeled by the momentum $\mathbf{k} = (k_x, k_y)$ and the irreducible representation of the axial point group. $(0,0)$ A $(S=0)$: (●), $(0,0)$ B $(S=0)$: (■), $(0,0)$ B $(S=1)$: (□), $(0,0)$ $x-iy$ $(S=1)$: (▽), $(0,\pi)$ A $(S=1)$: (+), (π,π) A $(S=1)$: (○), (π,π) $x+iy$ $(S=1)$: (△). Note that the ground state is in the ● sector.

To assess the validity of the hypotheses underlying Kodama *et al.*'s analysis beyond perturbation theory, we have performed exact diagonalizations on a 16 site cluster with periodic boundary conditions without making *a priori* assumptions on the value of the exchange integrals. First of all, we have determined the symmetry of the ground state and of the first triplet excitation in the (J_2/J_1,J_2'/J_1) plane for different values of J_1'/J_1 (see Fig. 1 for $J_1' = J_1$). Surprisingly enough, it turns out that only rather small values of J_2'/J_1 are compatible with the location of the minimum of the triplet band at $\mathbf{k} = (0, 0)$. To check the robustness of this result with respect to variations of the ratio J_1'/J_1, we have done a similar calculation as a function of (J_1'/J_1,J_2'/J_1) for $J_2/J_1 = 2$ (see Fig. 2). It shows that the restrictions are qualitatively the same, and that J_2'/J_1 must always be much smaller than J_2/J_1 in order to reproduce the neutron scattering results.



FIG. 2. Same as Fig. 1 for $J_2/J_1 = 2$.

Next, we have checked whether the set of parameters used in Ref. [2] and [3] was consistent with the picture of weakly coupled J_2-plaquettes. To answer this question we have calculated the spin-spin correlation function $\langle \mathbf{S}_i.\mathbf{S}_j \rangle$ with $J_1' = J_1$ in the presumed range of values of J_2/J_1 and J_2'/J_1. Fig. 3 summarizes the results and clearly delimits three different regions in which the system is organized into small J_1-plaquettes, large J_2-plaquettes and $J_1' - J_2'$ chains. We also checked that for $J_2/J_1 \gtrsim 2$ the correlation functions on all bonds except J_2 never exceed 0.1 in absolute value, which shows that the coupling between J_2-plaquettes is indeed very small in that range.



FIG. 3. "Phase diagram" in the (J_2/J_1, J_2'/J_1) plane. The symbols denote points where $\langle \mathbf{S}_i.\mathbf{S}_j \rangle < -0.4$ for J_1 (●),J_1' (○),J_2 (■) and J_2' (□) bonds.

In conclusion, the results obtained by a systematic study of the energy spectrum and of the spin-spin correlation functions of a 16 site cluster in a large range of J_1'/J_1,J_2/J_1 and J_2'/J_1 ratios show that another gapped phase made of weakly coupled J_2-plaquettes indeed exists when J_2/J_1 is large enough. Besides, the experimental fact that the minimum of the triplet band is located at $\mathbf{k} = (0, 0)$ implies that J_2' is much smaller than J_2. This suggests that the hypotheses of Kodama *et al.* are indeed satisfied. Calculations on a 32 site cluster to confirm the present results are in progress.

We thank IDRIS (Orsay) for allocation of CPU time on the C94 and C98 CRAY supercomputers.

---

[1] S. Taniguchi, T. Nishikawa, Y. Yashui, Y. Kobayashi, M. Sato, T. Nishioka, M. Kontani, and K. Sano. J. Phys. Soc. Jpn. **64**, 2758 (1995).

[2] K. Kodama, H. Harashina, H. Sasaki, Y. Kobayashi, M. Kasai, S. Taniguchi, Y. Yashui, M. Sato, K. Kakurai, K. Kakurai, and M. Nishi. J. Phys. Soc. Jpn. **66**, 28 (1997).

[3] K .Takano and K. Sano: cond-mat/9805153.